



## PATIENT INFORMATION LEAFLET (PIL)

**HemoMex S™**

**Softgel of Dammarane Sapogenins  
80 mg/Capsule**

### **WHAT YOU SHOULD KNOW ABOUT HemoMex S™ Softgel of Dammarane Sapogenins 80mg/Capsule**

Please read this leaflet carefully before you start to take HemoMex S™ Softgel of Dammarane Sapogenins 80mg/Capsule. It provides some information about your medicine and how to take it.

If you have any questions or are in any doubt about anything, please ask your doctor or pharmacist.

### **YOUR MEDICINE**

The name of your medicine is HemoMex S™ Softgel of Dammarane Sapogenins 80mg/Capsule. HemoMex S™ Softgel Capsule is a light brown, oval softgel capsule with light yellowish, oily content, astringent and bitter in taste.

### **WHAT IS IN YOUR MEDICINE?**

Each softgel capsule contains 80 mg Dammarane Sapogenins. It also contains Ethanol, Medium Chain Triglycerides, Polyoxyl 40 Hydrogenated Castor Oil, Gelatin, Glycerin, and Purified Water.

### **MANUFACTURER AND PRODUCT LICENCE HOLDER:**

Pegasus Pharmaceuticals Group Inc.  
Unit 110-116, 11800 River Road,  
Richmond, V6X 1Z7, Canada

### **WHAT IS YOUR MEDICINE FOR?**

HemoMex S™ Softgel Capsules are for:

Used in combination with chemotherapeutic agents and/or radiotherapy for solid tumors.

### **BEFORE YOU TAKE YOUR MEDICINE**

- Consult your physician prior to use or for use beyond 3 months. Do not use if you are pregnant or breast feeding.
- Consult your physician if you have any bleeding conditions, insomnia or schizophrenia, or you are taking any anticoagulant/antiplatelet

drugs, immunosuppressant drugs, steroids, antidepressant therapy, monoamine oxidase inhibitors (MAOI), and stimulant drugs.

- Patients taking warfarin (or other coumarin anticoagulants) should have increased monitoring of their INR levels when starting or stopping treatment with ginseng containing products.

### **HOW DO YOU TAKE YOUR MEDICINE?**

Take 5 capsules three times daily at 1-hour interval, 1 hour after lunch, for 5 consecutive days; repeat this regimen after a 2-day rest. Depending on individual's situation, may take continuously for up to twelve weeks or longer.

### **AFTER YOU HAVE TAKEN YOUR MEDICINE**

Normally this medicine is well tolerated but occasionally some people may get side effects.

Cases of headache, dizziness, insomnia, gastrointestinal reactions (such as nausea, stomach pain, vomiting and diarrhoea) and hypersensitivity reactions (such as rash, pruritus) have been observed.

### **HOW YOU SHOULD LOOK AFTER YOUR MEDICINE**

Store in a dry place. Do not freeze.

Keep out of reach of children.

Do not use this medicine after the expiry date which you will find printed on the packaging.

### **Distributed by:**

Pegasus Georgia Inc.  
Ap. 59, entrance 4, building 6, block 2, Gamsakhurdia Avenue  
Tbilisi, Georgia 0160  
Tel: +995 32 37 40 16

### **Further information is available to the medical and allied profession from:**

Pegasus Pharmaceuticals Group Inc.  
Unit 110-116, 11800 River Road,  
Richmond, V6X 1Z7, Canada  
Tel: +1-604-303-9988

**DRUG LICENCE NUMBER: No.R - 001177**

**DATE:** September 28, 2009